arXiv:1004.5390v1 [cond-mat.quant-gas] 29 Apr 2010

S-matrix approach to quantum gases in the unitary limit II: the three-dimensional case

Pye-Ton How and André LeClair
Newman Laboratory, Cornell University, Ithaca, NY

Abstract

A new analytic treatment of three-dimensional homogeneous Bose and Fermi gases in the unitary limit of negative infinite scattering length is presented, based on the S-matrix approach to statistical mechanics we recently developed. The unitary limit occurs at a fixed point of the renormalization group with dynamical exponent $z = 2$ where the S-matrix equals -1. For fermions we find $T_c/T_F \approx 0.1$. For bosons we present evidence that the gas does not collapse, but rather has a critical point that is a strongly interacting form of Bose-Einstein condensation. This bosonic critical point occurs at $n\lambda_T^3 \approx 1.3$ where n is the density and λ_T the thermal wavelength, which is lower than the ideal gas value of $\zeta(3/2) = 2.61$.

I. INTRODUCTION

In the so-called unitary limit of a quantum Bose or Fermi gas, the scattering length a diverges. This occurs at a fixed point of the renormalization group, thus these systems provide interesting examples of interacting, scale-invariant theories with dynamical exponent $z = 2$, i.e. non-relativistic. They can be realized experimentally by tuning the scattering length to $\pm\infty$ using a Feshbach resonance. (See for instance [1, 2] and references therein.) They are also thought to occur at the surface of neutron stars. These systems have also attracted much theoretical interest[3–17]. There have even been some proposals to use the AdS/CFT correspondence to learn about these models[18–21].

Because of the scale-invariance, the only length scales in the problem are based on the density $n^{1/d}$ where d is the spatial dimension, and the thermal wavelength $\lambda_T = \sqrt{2\pi/mT}$. Equivalently, the only energy scales are the chemical potential μ and the temperature T. The problem is challenging since there is no small paramater to expand in such as na^3. Any possible critical point must occur at a specific value of $x = \mu/T$. This can be translated into universal values for $n_c\lambda_T^3$, or for fermions universal values for T_c/T_F where $\epsilon_F = k_B T_F$ is the Fermi energy. For instance the critical point of an ideal Bose gas is the simplest example, where $n_c\lambda_T^3 = \zeta(3/2) = 2.61$.

The present work is the sequel to [22], where we used the S-matrix based formulation of the quantum statistical mechanics developed in[23, 24]. This approach is very well-suited to the problem because in the unitary limit the S-matrix $S = -1$, and kernels in the integral equations simplify. In fact, this approach can be used to develop an expansion in $1/a$. The main formulas for the 2 and 3 dimensional cases of both bosons and fermions were presented, however only the 2-dimensional case was analyzed in detail in [22]. Here we analyze the 3-dimensional case.

The models considered are the simplest models of non-relativistic bosons or fermions with quartic interactions. The bosonic model is defined by the action for a complex scalar field ϕ.

$$S = \int d^3\mathbf{x} dt \left(i\phi^\dagger \partial_t \phi - \frac{|\vec{\nabla}\phi|^2}{2m} - \frac{g}{4}(\phi^\dagger\phi)^2 \right) \tag{1}$$

For fermions, due to the fermionic statistics, one needs at least a 2-component field $\psi_{\uparrow,\downarrow}$:

$$S = \int d^3\mathbf{x} dt \left(\sum_{\alpha=\uparrow,\downarrow} i\psi_\alpha^\dagger \partial_t \psi_\alpha - \frac{|\vec{\nabla}\psi_\alpha|^2}{2m} - \frac{g}{2}\psi_\uparrow^\dagger\psi_\uparrow\psi_\downarrow^\dagger\psi_\downarrow \right) \tag{2}$$

In both cases, positive g corresponds to repulsive interactions. The bosonic theory only has a $U(1)$ symmetry. The fermionic theory on the other hand has the much larger SO(5) symmetry. This is evident from the work[32] which considered a relativistic version, since the same arguments apply to a non-relativistic kinetic term. This is also clear from the work[17] which considered an N-component version with Sp(2N) symmetry, and noting that Sp(4) = SO(5).

The interplay between the scattering length, the bound state, and the renormalization group fixed point was discussed in detail from the point of view of the S-matrix in [22]. In 3 spatial dimensions the fixed point occurs at negative coupling $g_* = -4\pi^2/m\Lambda$, where Λ is an ultra-violet cut-off. For g less than g_*, there is a bound state that can Bose-Einstein condense (BEC), and for the fermionic case, this is referred to as the BEC side. As g_* is approached from this side, the scattering length goes to $+\infty$. The bound state disappears at g_*. When g approaches g_* from above, the scattering length goes to $-\infty$, and this is referred to as the BCS side. In this paper we work on the BCS side of the cross-over since the bound state does not have to be incorporated into the thermodynamics. On this side the interactions are effectively attractive.

Theoretical studies have mainly focussed on the fermionic case, and for the most part at zero temperature, which is appropriate for a large Fermi energy. The bosonic case has been less studied, since a homogeneous bosonic gas with attractive interactions is thought to be unstable against mechanical collapse, and the collapse occurs before any kind of BEC. The situation is actually different for harmonically trapped gases, where BEC can occur[25]. However studies of the homogeneous bosonic case were based on a small, negative scattering length[26–29], and it is not clear that the conclusions reached there can be extrapolated to the unitary limit. Since the density of collapse is proportional to $1/a$[27], extrapolation to infinite scattering length suggests that the gas collapses at zero density, which seems unphysical, since the gas could in principle be stabilized at finite temperature by thermal pressure. One can also point out that in the van der Waals gas, the collapse is stabilized by a finite size of the atoms, which renders the compressibility finite. In the unitary limit, there is nothing to play such a role. In the sequel we will present evidence that the unitary Bose gas undergoes BEC when $n\lambda_T^3 \approx 1.3$. This lower value is consistent with the attractive interactions. We also estimate the critical exponent describing how the compressibility diverges at the critical point.

In the next section we define the scaling functions that determine the free energy and density, and derive expressions for the energy and entropy per particle, specific heat per particle, and compressibility. In section III the formulation of the unitary limit in [22] is summarized. The two-component fermion case is analyzed in section IV. Evidence is presented for a critical point with $T_c/T_F \approx 0.1$, which is consistent with lattice Monte Carlo simulations. Bosons are analyzed in section V, where we present evidence for BEC in this strongly interacting gas. Motivated by the conjectured lower bound[30] for the ratio of the viscosity to entropy density $\eta/s > \hbar/4\pi k_B$ for relativistic systems, we study this ratio for both the fermionic and bosonic cases in section VI. Our results for fermions are consistent with experiments, with $\eta/s > 4.72$ times the conjectured lower bound. For bosons, this ratio is minimized at the critical point where $\eta/s > 1.26$ times the bound.

II. SCALING FUNCTIONS IN THE UNITARY LIMIT.

The scale invariance in the unitary limit implies some universal scaling forms[6]. In this section we define various scaling functions with a meaningful normalization relative to free particles.

First consider a single species of bosonic or fermionic particle with mass m at chemical potential μ and temperature T. The free energy density has the form

$$\mathcal{F} = -\zeta(5/2)\left(\frac{mT}{2\pi}\right)^{3/2} T\, c(\mu/T) \tag{3}$$

where the scaling function c is only a function of $x \equiv \mu/T$. (ζ is Riemann's zeta function.) The combination $\sqrt{mT/2\pi} = 1/\lambda_T$, where λ_T is the thermal wavelength. For a single free boson or fermion:

$$\lim_{\mu/T\to 0} \quad c_{\rm boson} = 1, \qquad c_{\rm fermion} = 1 - \frac{1}{2\sqrt{2}} \qquad \text{(free particles)}. \tag{4}$$

It is also convenient to define the scaling function q, which is a measure of the quantum degeneracy, in terms of the density as follows:

$$n\lambda_T^3 = q \tag{5}$$

The two scaling functions c and q are of course related since $n = -\partial\mathcal{F}/\partial\mu$, which leads to

$$q = \zeta(5/2)c' \tag{6}$$

where c' is the derivative of c with respect to x. Henceforth g' will always denote the derivative of g with respect to x. The expressions for c and q for free theories will be implicit in the next section.

Also of interest are several energy per particle scaling functions. At a renormalization group fixed point, the energy density is related to the free energy in the same way as for free particles:

$$\frac{E}{V} = -\frac{3}{2}\mathcal{F} \tag{7}$$

where V is the volume. For a free fermion, in the zero temperature limit, the energy per particle $E/N \to \frac{3}{5}\mu$. The Fermi energy is

$$\epsilon_F = \frac{1}{m}\left(3\pi^2 n/\sqrt{2}\right)^{2/3} \tag{8}$$

The above definition can also be used for bosons. Since $\epsilon_F = \mu$ in the zero temperature free fermionic gas, this leads us to define the scaling function ξ:

$$\xi(x) = \frac{5}{3}\frac{E}{N\epsilon_F} = \frac{5\zeta(5/2)}{3}\left(\frac{6}{\pi}\right)^{1/3}\frac{c}{q^{5/3}} \tag{9}$$

In the limit $T \to 0$, i.e. $x \to \infty$, $\xi \to 1$ for a free fermion.

A different energy per particle scaling function, $\widetilde{\xi}$, is meaningful as $x \to 0$:

$$\frac{E}{NT} = \frac{3(2\sqrt{2}-1)\zeta(5/2)}{2(2\sqrt{2}-2)\zeta(3/2)}\,\widetilde{\xi}(x) \tag{10}$$

With the above normalization $\widetilde{\xi} = 1$ for a free fermion in the limit $x \to 0$. In terms of the above scaling functions:

$$\widetilde{\xi}(x) = \frac{\zeta(3/2)(2\sqrt{2}-2)}{(2\sqrt{2}-1)}\,\frac{c}{q} \tag{11}$$

The entropy density is $s = -\partial\mathcal{F}/\partial T$, and the entropy per particle takes the form

$$\frac{s}{n} = \zeta(5/2)\left(\frac{5c/2 - xc'}{q}\right) \tag{12}$$

Next consider the specific heat per particle at constant volume and particle number, i.e. constant density. One needs $\partial x/\partial T$ at constant density. Using the fact that $n \propto T^{3/2}q$, at constant density $q \propto T^{-3/2}$. This gives

$$T\left(\frac{\partial x}{\partial T}\right)_n = -\frac{3}{2}\frac{q}{q'} \tag{13}$$

The specific heat per particle is then:

$$\frac{C_V}{N} = \frac{1}{N}\left(\frac{\partial E}{\partial T}\right)_{N,V} = \frac{\zeta(5/2)}{4}\left(15\frac{c}{q} - 9\frac{c'}{q'}\right) \tag{14}$$

The isothermal compressibility is defined as

$$\kappa = -\frac{1}{V}\left(\frac{\partial V}{\partial p}\right)_T \tag{15}$$

where the pressure $p = -\mathcal{F}$. Since $n = N/V$ and N is kept fixed,

$$\kappa = -n\left(\frac{\partial n^{-1}}{\partial p}\right)_T = \frac{1}{nT}\frac{q'}{q} = \frac{1}{T}\left(\frac{mT}{2\pi}\right)^{3/2}\frac{q'}{q^2} \tag{16}$$

Finally the equation of state can be expressed parametrically as follows. Given n and T, one uses eq. (5) to find x as a function of n, T. The pressure can then be written as

$$p = \left(\frac{\zeta(5/2)c(x(n,T))}{q(x(n,T))}\right) nT \tag{17}$$

In order to compare with numerical simulations and experiments, it will be useful to plot various quantities as a function of q or T/T_F:

$$\frac{T}{T_F} = \left(\frac{4}{3\sqrt{\pi}q}\right)^{2/3} \tag{18}$$

III. TWO-BODY SCATTERING APPROXIMATION

The main features of the two-body scattering approximation developed in [24] are the following. Consider again first a single component gas. The filling fractions, or occupation numbers, are parameterized in terms of a pseudo-energy $\varepsilon(\mathbf{k})$:

$$f(\mathbf{k}) = \frac{1}{e^{\beta\varepsilon(\mathbf{k})} - s} \tag{19}$$

which determine the density:

$$n = \int \frac{d^3\mathbf{k}}{(2\pi)^3}\,\frac{1}{e^{\beta\varepsilon(\mathbf{k})} - s} \tag{20}$$

where $s = 1, -1$ corresponds to bosons, fermions respectively and $\beta = 1/T$. The consistent summation of 2-body scattering leads to an integral equation for the pseudo-energy $\varepsilon(\mathbf{k})$. It is convenient to define the quantity:

$$y(\mathbf{k}) = e^{-\beta(\varepsilon(\mathbf{k}) - \omega_{\mathbf{k}} + \mu)} \tag{21}$$

where $\omega_{\mathbf{k}} = \mathbf{k}^2/2m$. Then y satisfies the integral equation

$$y(\mathbf{k}) = 1 + \beta \int \frac{d^3\mathbf{k}'}{(2\pi)^3} G(\mathbf{k} - \mathbf{k}') \frac{y(\mathbf{k}')^{-1}}{e^{\beta\varepsilon(\mathbf{k}')} - s} \tag{22}$$

The free energy density is then

$$\mathcal{F} = -T \int \frac{d^3\mathbf{k}}{(2\pi)^3} \left[-s \log(1 - s e^{-\beta\varepsilon}) - \frac{1}{2} \frac{(1 - y^{-1})}{e^{\beta\varepsilon} - s} \right] \tag{23}$$

The kernel has the following structure:

$$G = -\frac{i}{\mathcal{I}} \log(1 + i\mathcal{I}\mathcal{M}) \tag{24}$$

where $\mathcal{M}$ is the scattering amplitude and $\mathcal{I}$ represents the available phase space for two-body scattering. The argument of the log can be identified as the S-matrix function. In the unitary limit,

$$\mathcal{M} = \frac{2i}{\mathcal{I}} = \frac{16\pi i}{m|\mathbf{k} - \mathbf{k}'|}, \tag{25}$$

and the S-matrix equals -1. The kernel becomes

$$G(\mathbf{k} - \mathbf{k}') = \mp \frac{8\pi^2}{m|\mathbf{k} - \mathbf{k}'|}, \tag{26}$$

where the $-$ sign corresponds to g being just below the fixed point g_*, where the scattering length $a \to +\infty$ on the BEC side, whereas the $+$ sign corresponds to $a \to -\infty$ on the BCS side. As explained in the Introduction, we work on the BCS side.

The angular integrals in eq. (22) are easily performed. Defining the dimensionless variable $\kappa = \mathbf{k}^2/2mT$, the integral equation becomes

$$y(\kappa) = 1 + 4\int_0^\infty d\kappa' \left[\Theta(\kappa-\kappa')\sqrt{\kappa'/\kappa} + \Theta(\kappa'-\kappa)\right] \frac{z}{e^{\kappa'} - szy(\kappa')} \tag{27}$$

where $z = e^{\mu/T}$ is the fugacity and $\Theta(\kappa)$ is the standard step function equal to 1 for $\kappa > 0$, zero otherwise.

Finally comparing with the definitions in the last section the scaling function for the density and free energy are

$$q(x) = \frac{2}{\sqrt{\pi}}\int_0^\infty d\kappa\sqrt{\kappa}\frac{y(\kappa)z}{e^\kappa - sy(\kappa)z} \tag{28}$$

and

$$c = \frac{2}{\sqrt{\pi}\zeta(5/2)}\int_0^\infty d\kappa\sqrt{\kappa}\left(-s\log\left(1 - szy(\kappa)e^{-\kappa}\right) - \frac{1}{2}\frac{z(y(\kappa)-1)}{e^\kappa - szy(\kappa)}\right) \tag{29}$$

The ideal, free gas limit corresponds to $y = 1$ where $q = s\mathrm{Li}_{3/2}(sz)$ and $c = s\mathrm{Li}_{5/2}(sz)/\zeta(5/2)$, where Li is the polylogarithm. The BEC critical point of the ideal gas occurs at $\mu = 0$, i.e. $q = \zeta(3/2)$.

Consider now two-component fermions with the action (2). Here the phase space factor $\mathcal{I}$ is doubled and since $G \propto 1/\mathcal{I}$, the kernels have an extra 1/2:

$$G_{\text{fermi}} = \frac{1}{2}G_{\text{bose}} \tag{30}$$

Due to the SU(2) symmetry, the two-component fermion reduces to two identical copies of the above 1-component expressions, with the modification (30).

IV. ANALYSIS OF FERMIONS

Recall that for 2 component fermions, the 4 is replaced by 2 in eq. (27), and by the SU(2) symmetry, the system reduces to two identical copies of the one-component equations of the last sections; below we present results for a single component, it being implicit that the free energy and density are doubled.

The integral equation for $y(\kappa)$, eq. (27), can be solved numerically by iteration. One first substitutes $y_0 = 1$ on the right hand side and this gives the approximation y_1 for y. One then substitutes y_1 on the right hand side to generate y_2, etc. For regions of z where there are no critical points, this procedure converges rapidly, and as little as 5 iterations are needed. For fermions, as one approaches zero temperature, i.e. x large and positive, more iterations are needed for convergence. The following results are based on 50 iterations.

When $z \ll 1$, $y \approx 1$, and the properties of the free ideal gas are recovered, since the gas is very dilute. There are solutions to eq. (27) for all $-\infty < x < \infty$. ($x = \mu/T$). The scaling function c, and it's comparision with a free theory, are shown in Figure 1 as a function of x. The corrections to the free theory become appreciable when $x > -2$. At $x = 0$:

$$c(0) = 0.880, \qquad \widetilde{\xi} = 0.884, \tag{31}$$

compared to the free gas values of $c(0) = 0.646$ and $\widetilde{\xi} = 1$.

The scaling function q for the density is shown as function of x in Figure 2. Note that the density in the interacting case is always higher than for a free gas, due to the attractive interactions. At $x = 0$, $q(0) = 1.18$, whereas for a free gas $q = 0.765$. At low temperatures and high densities, $\mu/T \gg 1$, the occupation numbers resemble that of a degenerate Fermi gas, as shown in Figure 3.

Whereas c and q are nearly featureless, other quantities seem to indicate a critical point, or phase transition, at large density. For instance, the entropy per particle decreases with decreasing temperature up to $x < x_c \approx 11.2$, as shown in Figure 4. Beyond this point the entropy per particle has the unphysical behavior of increasing with temperature. A further indication that the region $x > x_c$ is unphysical is that



FIG. 1: $c(x)$ and its equivalent for a free theory as a function of $x = \mu/T$.

the specific heat per particle becomes negative, as shown in Figure 5. When $x \ll 0$, C_V/N approaches the classical value 3/2. This leads us to suggest a phase transition, at $x = x_c$, corresponding to the critical temperature $T_c/T_F \approx 0.1$. As we will show, our analysis of the viscosity to entropy-density ratio suggests a higher T_c/T_F. There have been numerous estimates of T_c/T_F based on various approximation schemes, mainly using Monte Carlo methods on the lattice [10–15], quoting results for T_c/T_F between 0.05 and 0.23. The work [11] puts an upper bound $T_c/T_F < 0.14$, and the most recent results of Burovski et. al. quote $T_c/T_F = 0.152(7)$. Our result is thus consistent with previous work. The equation of state at this point follows from eq. (17):

$$p = 4.95nT \tag{32}$$

The energy per particle, normalized to the Fermi energy ϵ_F, i.e. $E/N\epsilon_F = 3\xi/5$, and the entropy per particle, are shown in Figures 6,7 as a function of T/T_F, where $k_B T_F = \epsilon_F$. At high temperatures it matches that of a free Fermi gas, in agreement



FIG. 2: $q(x)$ and its equivalent for a free theory as a function of $x = \mu/T$.

with the Monte Carlo simulations in [13, 15]. Note that there is no sign of pair-breaking at $T^*/T_F = 0.5$ predicted in [10], and this also agrees with the Monte Carlo simulations. However at low temperatures in the vicinity of T_c, the agreement is not as good. This suggests our approximation is breaking down for very large z, i.e. the limit of zero temperature. The same conclusion is reached by examining μ/ϵ_F, displayed in Figure 8, since the zero temperature epsilon expansion and Monte Carlo give $\mu/\epsilon_F \approx 0.4 - 0.5$[15, 18].

V. ANALYSIS OF BOSONS

For bosons we again solved the integral equation (27) by iteration, starting from $y = 1$. Since the occupation numbers decay quickly as a function of κ, we introduced a cut-off $\kappa < 10$. For x less than approximately -2, the gas behaves nearly classically.

The main feature of the solution to the integral equation is that for $x > x_c \equiv$



FIG. 3: The occupation numbers as a function of κ for $x = 5, 10, 15$.

-1.2741, there is no solution that is smoothly connected to the classical limit $x \to -\infty$. Numerically, when there is no solution the iterative procedure fails to converge. The free energy scaling function is plotted in Figure 9. Note that $c < 1$, where $c = 1$ is the free field value. We thus take the physical region to be $x < x_c$. We find strong evidence that the gas undergoes BEC at $x = x_c$. In Figure 10, we plot $\varepsilon(\mathbf{k} = 0)$ as a function of x, and ones sees that it goes to zero at x_c. This implies the occupation number f diverges at $\mathbf{k} = 0$ at this critical point. One clearly sees this behavior in Figure 11.

The compressibility is shown in Figure 12, and diverges at x_c, again consistent with BEC. We thus conclude that there is a critical point at x_c which a strongly interacting, scale invariant version of the ideal BEC. In terms of the density, the critical point is:

$$n_c \lambda_T^3 = 1.325, \qquad (\mu/T = x_c = -1.2741) \tag{33}$$



FIG. 4: Entropy per fermionic particle as a function of x.

The negative value of the chemical potential is consistent with the effectively attractive interactions. The above should be compared with the ideal BEC of the free theory, where $x_c = 0$ and $n_c \lambda_T^3 = \zeta(3/2) = 2.61$, which is higher by a factor of 2. At the critical point the equation of state is

$$p = 0.318 nT \tag{34}$$

compared to $p = 0.514 nT$ for the free case. $(0.514 = \zeta(5/2)/\zeta(3/2))$.

A critical exponent ν characterizing the diverging compressibility can be defined as

$$\kappa \sim (T - T_c)^{-\nu} \tag{35}$$

A log-log plot of the compressibility verses $T - T_c$ shows an approximately straight line, and we obtain $\nu \approx 0.69$. This should be compared with BEC in an ideal gas, where $\nu \approx 1.0$. Clearly the unitary gas version of BEC is in a different universality class.



FIG. 5: Specific heat per particle as a function of x for fermions.



FIG. 6: Energy per particle normalized to ϵ_F as a function of T/T_F.

The energy per particle scaling function $\widetilde{\xi}$ at the critical point is $\widetilde{\xi}(x_c) = 0.281$ compared to $0.453 = (2\sqrt{2}-2)/(2\sqrt{2}-1)$ for the free case. The entropy per particle and specific heat per particle are plotted in Figures 13, 14 as a function of T/T_c. At large temperatures, as expected $C_V/N = 3/2$, i.e. the classical value. It increases as



FIG. 7: Entopy per particle as a function of T/T_F.



FIG. 8: Chemical potential normalized to ϵ_F as a function of T/T_F.

T is lowered, however in contrast to the ideal gas case, it then begins to decrease as T approaches T_c.



FIG. 9: The free-energy scaling function c as a function of μ/T compared to the ideal gas case.

VI. ENTROPY TO VISCOSITY RATIO

Consider first a single component gas. In kinetic theory, the shear viscosity can be expressed as

$$\eta = \frac{1}{3} n \overline{v} m \ell_{\text{free}} \tag{36}$$

where $\overline{v}$ is the average speed and ℓ_{free} is the mean free path. The mean free path is $\ell_{\text{free}} = 1/(\sqrt{2} n \sigma)$ where σ is the total cross-section. (The $\sqrt{2}$ comes from the ratio of the mean speed to the mean relative speed[34].) In the unitary limit the S-matrix $S = -1$, which implies the scattering amplitude in eq. (25). This leads to

$$\sigma = \frac{m^2 |\mathcal{M}|^2}{4\pi} = \frac{16\pi}{|\mathbf{k}|^2} \tag{37}$$

where $|k|$ is the momentum of one of the particles in the center of mass frame, i.e. $|\mathbf{k}_1 - \mathbf{k}_2| = 2|\mathbf{k}|$. This gives

$$\eta = \frac{m^3 \overline{v}^3}{48\sqrt{2}\pi} \tag{38}$$



FIG. 10: The pseudo-energy ε at $\mathbf{k} = 0$ as a function of $x = \mu/T$.

Since the equation (7) is the same relation between the pressure and energy of a free gas, and the pressure is due to the kinetic energy, this implies

$$\frac{1}{2} m \overline{v}^2 = E/N = \frac{3}{2} \frac{c}{c'} T \tag{39}$$

Since the entropy density $s = -\partial \mathcal{F}/\partial T$, one finally has

$$\frac{\eta}{s} = \frac{\sqrt{3\pi}}{8\zeta(5/2)} \left(\frac{c}{c'}\right)^{3/2} \frac{1}{5c/2 - xc'} \tag{40}$$

For two-component fermions, the available phase space $\mathcal{I}$ is doubled. Also, spin up particles only scatter with spin down. This implies η is 8 times the above expression. Since the entropy density is doubled, this implies that η/s is 4 times the expression eq. (40).

The ratio η/s for fermions as a function of T/T_F is shown in Figure 15, and is in good agreement both quantitatively and qualitatively with the experimental data summarized in [37]. The lowest value occurs at $x = 2.33$, which corresponds to



FIG. 11: The occupation number $f(\kappa)$ for $x = -1.275$ and $x_c = -1.2741$.

$T/T_F = 0.28$, and

$$\frac{\eta}{s} > 4.72 \frac{\hbar}{4\pi k_B} \tag{41}$$

The experimental data has a minimum that is about 6 times this bound. In the free fermion theory the minimum occurs at $\mu/T \approx 2.3$, which gives $\eta/s > 7.2\hbar/4\pi k_B$.

For bosons, the ration η/s is plotted in Figure 16 as a function of T/T_c. One sees that it has a minimum at the critical point, where

$$\frac{\eta}{s} > 1.26 \frac{\hbar}{4\pi k_B} \tag{42}$$

Thus the bosonic gas at the unitary critical point is a more perfect fluid than that of fermions. On the other hand, the ideal Bose gas at the critical point has a lower value:

$$\left.\frac{\eta}{s}\right|_{\text{ideal}} = \frac{\sqrt{3\pi\zeta(5/2)}}{20\zeta(3/2)^{3/2}} = 0.53 \frac{\hbar}{4\pi k_B} \tag{43}$$



FIG. 12: The compressibility κ as a function of μ/T.



FIG. 13: The entropy per particle as a function of T/T_c.



FIG. 14: The specific heat per particle as a function of T/T_c.



FIG. 15: The viscosity to entropy-density ratio as a function of T/T_F for fermions. The horizontal line is $1/4\pi$.

VII. CONCLUSIONS

We presented a novel analytic treatment of unitary Bose and Fermi gases at finite temperature and chemical potential using a new formulation of statistical mechanics



FIG. 16: The viscosity to entropy-density ratio as a function of T/T_c for bosons. The horizontal line is $1/4\pi$.

based on the exact, zero temperature, 2-body scattering. Our results appear to be consistent with lattice Monte Carlo methods. All of the thermodynamic functions, such as entropy per particle, energy per particle, specific heat, compressibility, and viscosity are readily calculated once one numerically solves the integral equation for the pseudo-energy.

For fermions, our 2-body approximation is good if the temperatures are not too low. We estimated $T_c/T_F \approx 0.1$, where the critical point occurs at $\mu/T \approx 11.2$. For bosons we presented evidence for a strongly interacting version of BEC at the critical point $n\lambda_T^3 \approx 1.3$, corresponding to $\mu/T = -1.27$.

VIII. ACKNOWLEDGMENTS

We wish to thank Erich Mueller for helpful discussions. This work is supported by the National Science Foundation under grant number NSF-PHY-0757868.

---

[1] K. M. O'Hara et. al., Science **298** (2002) 2179.

[2] J. Kinast et. al. Science **298** (2002) 2179.

[3] A. J. Leggett, in *Modern Trends in the Theory of Condensed Matter*, Springer, Berlin 1980.

[4] P. Nozières and S. Schmitt-Rink, J. Low Temp. Phys. **59** (1985) 195.

[5] Y. Ohashi and A. Griffin, *The BCS-BEC Crossover in a Gas of Fermi Atoms with a Feshbach Resonance*, Phys. Rev. Lett. **89** (2002) 130402 [arXiv:cond-mat/0210185].

[6] T.-L. Ho, *Universal Thermodynamics of Degenerate Quantum Gases in the Unitary Limit*, Phys. Rev. Lett. **92** (2004) 090402 [arXiv:cond-mat/0309109].

[7] T.-L. Ho and E. Mueller, *High temperature expansion applied to fermions near a Feshbach resonance*, Phys.Rev.Lett. **92** (2004) 160404 [arXiv:cond-mat/0306187].

[8] G.E. Astrakharchik, J. Boronat, J. Casulleras and S. Giorgini, *Equation of state of a Fermi gas in the BEC-BCS crossover: a quantum Monte Carlo study*, Phys. Rev. Lett. **93** (2004) 200404 [arXiv:cond-mat/0406113].

[9] Z. Nussinov and S. Nussinov, *The BCS-BEC Crossover in Arbitrary Dimensions*, [arXiv:cond-mat/0609106].

[10] A. Perali, P. Pieri, L. Pisani and G. C. Strinati, *BCS-BEC crossover at finite temperature for superfluid trapped Fermi atoms*, Phys. Rev. Lett. **87** (2004) 220404 [cond-mat/0311309].

[11] D. Lee and T. Schäfer, *Cold dilute neutron matter on the lattice II: Results in the*

unitary limit, Phys.Rev. **C73** (2006) 015202 [nucl-th/0509018].

[12] M. Wingate, *Critical temperature for fermion pairing using lattice field theory*, [cond-mat/0502372]

[13] A. Bulgac, J. E. Drut and P. Magierski, *Spin 1/2 Fermions in the Unitary Regime: A Superfluid of a New Type*, Phys. Rev. Lett. **96** (2006) 090404 [arXiv:cond-mat/050537]; *Spin 1/2 Fermions in the Unitary Regime at Finite Temperature*, Int. J. Mod. Phys. **B20**, 5165 (2006) [arXiv:cond-mat/0602091].

[14] H. Hu, X.J. Liu and P. D. Drummond, *Equation of state of a superfluid Fermi gas in the BCS-BEC crossover*, Europhys. Lett. **74**(2006) 574 [cond-mat/0506046].

[15] E. Burovski, N. Prokof'ev, B. Svistunov and M. Troyer, *The Fermi-Hubbard model at unitarity*, New J.Phys.**8** (2006) 153 [arXiv:cond-mat/0605350]; *Critical Temperature and Thermodynamics of Attractive Fermions at Unitarity*, Phys. Rev. Lett. **96** (2006) 160402.

[16] Y. Nishida and D. T. Son, ϵ expansion for a Fermi gas at infinite scattering length, Phys. Rev. Lett. **97** (2006) 050403 [arXiv:cond-mat/0604500].

[17] P. Nikolić and S. Sachdev, *Renormalization group fixed points, universal phase diagram, and 1/N expansion for quantum liquids with interactions near the unitarity limit*, Phys. Rev. **A75** (2007) 033608 [arXiv:cond-mat/0609106].

[18] D. T. Son, *Toward an AdS/cold atoms correspondence: a geometric realization of the Schoedinger symmetry*, Phys. Rev. **D78** (2008) 046003 [arXiv:0804.3972].

[19] J. Maldacena, D. Martelli and Y. Tachikawa, *Comments on string theory backgrounds with non-relativistic conformal symmetry*, JHEP0810 (2008) 072 [arXiv:0807.1100].

[20] C. P. Herzog, M. Rangamani and S. F. Ross, *Heating up Galilean holography*, JHEP0811 (2008) 080 [arXiv:0807.1099].

[21] A. Adams, K. Balasubramanian and J. McGreevy, *Hot spacetimes for cold atoms*, JHEP0811:059, 2008 [arXiv:0807.1111].

[22] P.-T. How and A. LeClair, *S-matrix approach to quantum gases in the unitary limit I: the two-dimensional case*, to appear JSTAT.

[23] A. LeClair, *Quantum statistical mechanics of gases in terms of dynamical filling fractions and scattering amplitudes*, J.Phys. **A40** (2007) 9655 [arXiv:hep-th/0611187].

[24] P.-T. How and A. LeClair, *Critical point of the two-dimensional Bose gas: an S-matrix approach*, Nucl. Phys. **B824** (2010) 415 [arXiv:0906.0333].

[25] C. C. Bradley, C. A. Sackett, J. J. Tollett, and R. G. Hulet, *Evidence of Bose-Einstein condensation in an atomic gas with attractive interactions*, Phys. Rev. Lett. **75** (1995) 1687.

[26] H. T. C. Stoof, *Atomic Bose gas with a negative scattering length*, Phys. Rev. **A49** 3824.

[27] E. J. Mueller and G. Baym, *Finite Temperature Collapse of a Bose Gas with Attractive Interactions*, [cond-mat/0005323]

[28] G. S. Jeon, L. Yin, S. W. Rhee and D. J. Thouless, *Pairing instability and Mechanical Collapse of a Bose Gas with an Attractive Interaction*, [cond-mat/0110004]

[29] Yu-Li Lee and Yu-Wen Lee, *The Universality and stability for a dilute Bose gas with a Feshbach resonance*, [arXiv:1001.2592].

[30] P. Kovtun, D. T. Son and A. O. Starinets, Phys. Rev. Lett. **94** (2005) 111606 [arXiv:hep-th/0405231].

[31] D. S. Petrov, M. A. Baranov and G. V. Shlyapnikov, *Superfluid transition in quasi 2d Fermi gases*, Phys. Rev. **A 67** (2003) 031601 [arXiv:cond-mat/02112061].

[32] E. Kapit and A. LeClair, *A model of a 2d non-Fermi liquid with SO(5) symmetry, AF order, and a d-wave SC gap*, J. Phys. **A42** (2009) 025402 [arXiv:0805.4182].

[33] A. LeClair, *Interacting Bose and Fermi gases in low dimensions and the Riemann hypothesis*, Int. J. Mod. Phys. **A23** (2008) 1371 [arXiv:math-ph/0611043].

[34] F. Reif, *Fundamentals of statistical and thermal physics*, McGraw-Hill 1965.

[35] P. Massignan, G. M. Bruun and H. Smith, *Viscous relaxation and collective oscillations in a trapped Fermi gas near the unitarity limit*, Phys. Rev. **A71** (2005) 033607 [arXiv:cond-mat/0409660].

[36] B. A. Gelman, E. V. Shuryak and I. Zahed, *Cold Strongly Coupled Atoms Make a Near-perfect Liquid*, Phys. Rev. **A72** (2005) 043601 [arXiv:nucl-th/0410067].

[37] T. Schäfer, *The Shear Viscosity to Entropy Density Ratio of Trapped Fermions in the Unitarity Limit,* Phys. Rev. **A76** (2007) 063618 [arXiv:cond-mat/0701251].

[38] G. Rupak and T. Schafer, *Shear viscosity of a superfluid Fermi gas in the unitarity limit*, Phys. Rev. **A76** (2007) 073607 [arXiv:0707,1520].

[39] R. S. Crandall and R. Williams, Phys. Lett. **34A** (1971) 404.

[40] C. C. Grimes and G. Adams, *Evidence for a Liquid-to-Crystal Phase Transition in a Classical, Two-Timensional Sheet of Electrons*, Phys. Rev. Lett. **42** (1979) 795.